Hay oat guide

SA
Vic
NSW

ARCHER$^{\Phi}$

ARCHER$^{\Phi}$ offers

- Single gene IMI tolerant hay variety, an ideal variety for use where there are IMI residue concerns from previous crops.
- Mid-maturity with a mid-tall plant height.
- Moderate stem diameter. Seeding density should be managed to ensure stem diameter does not get too thick and reduce water soluble carbohydrate.
- A good disease profile including excellent resistance to oat leaf (crown) rust.

imiCrops®

KINGBALE$^{\Phi}$

KINGBALE$^{\Phi}$ offers

- A single gene IMI tolerant oaten hay variety, an ideal variety for use where there are IMI residue concerns from previous crops.
- Mid-slow maturing and tall variety with good early vigour.
- Excellent CCN resistance (R), useful in paddocks with a tight cereal rotation.

imiCrops®

GOLDIE$^{\Phi}$

GOLDIE$^{\Phi}$ offers

- Mid maturity, 4-7 days earlier to flower than Bannister$^{\Phi}$
- A strong hay yield and hay quality as well as strong grain yields and quality package.
- Good CCN resistance (MR) and oat crown rust (RMR).
- A tall plant height and good panicle emergence even under tough conditions, an improvement compared to Bannister$^{\Phi}$.

WALLABY$^{\Phi}$

WALLABY$^{\Phi}$ offers

- A mid-slow maturing oaten hay variety for April sowing, approximately 7-10 days slower than Bannister$^{\Phi}$.
- Market leading quality attributes including good digestibility and high water-soluble carbohydrate levels.
- Medium to tall plant height and well likely suited to medium and high rainfall zones.
- Good CCN resistance.

Hay Oat Guide

SA
Vic
NSW

Hay yield and quality performance

	Hay Yield (t/ha)	NDF (% dm)	ADF (% dm)	NDFdom30	Crude Protein (% dm)	Digestability (% dm)	WSC (% dm)	Grain Yield (t/ha)
Archer[◊]	10.5	54.7	33.1	49.3	7.2	62.3	21.4	5.0
Kingbale[◊]	10.7	55.2	34.1	48.3	6.3	62.0	22.7	3.9
Goldie[◊]	10.8	53.5	32.0	50.4	6.8	64.4	24.6	5.2
Wallaby[◊]	10.6	53.4	31.7	50.8	6.7	65.4	25.1	4.6
Bannister[◊]	10.3	53.9	31.7	50.5	6.9	64.7	23.3	5.1
Brusher[◊]	10.4	54.1	32.6	49.1	6.9	63.6	23.6	3.9
Mulgara[◊]	10.3	54.2	33.0	49.7	6.5	63.3	24.0	3.9

NDF = Neutral detergent fibre, ADF = Acid detergent fibre, WSC = Water-soluble carbohydrates, NDFDom30 = neutral detergent fibre digestibility after 30 hours. Source: 2018-2025 InterGrain national MET. Goldie[◊] only present in 2021-25.

Disease

	Archer[◊]	Kingbale[◊]	Goldie[◊]	Wallaby[◊]
Septoria blotch	MSS	MS	MSS	MSS
Leaf (crown) rust (East)	MR-S	S	S	MR
Stem rust	MSS	S	S	SVS
Barley and cereal yellow dwarf virus	MSS	MS	MS	MS
Bacterial blight	MSS	MSS	MSS	S
CCN resistance	VS	R	MR	MR
Red leather leaf	SVS	SVS	SVS	SVS
P. neglectus resistance	MSp	MRMS	MSp	MRMS
P. thornei resistance	MRMS	MRMS	MRMS	MS
Stem nematode resistance	VSp	MR	S	S
Stem nematode tolerance	I	MT	I	MI

Source: 2026 NVT Pathology consensus disease ratings. p= provisional rating, R = Resistant, RMR = Resistant to Moderately Resistant, MR = Moderately Resistant, MRMS = Moderately Resistant to Moderately Susceptible, MS = Moderately Susceptible, MSS = Moderately Susceptible to Susceptible, S = Susceptible, SVS = Susceptible to Very Susceptible, VS = Very Susceptible, T = Tolerant, T-MT = Tolerant – Moderately Tolerant, MT = Moderately Tolerant, MT-MI = Moderately Tolerant – Moderately Intolerant, MI = Moderately Intolerant, MI-I = Moderately Intolerant – Intolerant, I = Intolerant, I-VI = I Intolerant – Very Intolerant, IV = Very Intolerant

Hay Oat Guide

SA
Vic
NSW

Disease

	Archer(ф)	Wallaby(ф)	Kingbale(ф)	Goldie(ф)
Variety type	Hay	Hay	Hay	Milling/ hay
Maturity	Mid	Mid-slow	Mid-slow	Mid
Plant height	Mid-tall	Mid-tall	Tall	Mid-tall
Coleoptile + mesocotyl length	>120mm	>120mm	>120mm	>120mm
Stem diameter	Moderate	Moderate-fine	Moderate-fine	Moderate-fine

Source: InterGrain Oat Breeding

IMI Oat Stewardship

KINGBALE(ф) and ARCHER(ф) are single gene IMI tolerant oaten hay varieties. Sentry® is registered for incorporation by seeding (IBS) for hay, forage, seed and grain (domestic feed market only) production.

Please note that excess grain, seed and screenings produced from KINGBALE(ф) and ACRHER(ф) can only be used within domestic oaten grain feed markets and/or consumed on-farm. Grain of these varieties cannot be delivered into export bulk handling supply chains.

KINGBALE(ф) and ARCHER(ф) are available through resellers, Seedclub Members Farmer-to-Farmer Trade is not permitted.

The original breeding work for KINGBALE(ф) and ACRHER(ф) was undertaken by Grains Innovation Australia (GIA). InterGrain is commercializing the varieties.

GRAINS INNOVATION AUSTRALIA

Nufarm

InterGrain Territory Manager contact

Darcy Moore
South Australia
0448 138 944 | dmoore@intergrain.com

Dan Fay
Victoria
0427 850 766 | dfay@intergrain.com

David Leah
Southern New South Wales
0447 565 457 | dleah@intergrain.com

Matt Naumann
Northern New South Wales & Queensland
0460 292 620 | mnaumann@intergrain.com

PBR/EPR
InterGrain varieties are protected by Plant Breeder's Rights and are subject to an end point royalty for grain and hay production.
ARCHER(ф) and **KINGBALE**(ф) $3.65/tonne GST exclusive
WALLABY(ф) $3.00/tonne GST exclusive
GOLDIE(ф) $3.50/tonne GST exclusive

Disclaimer
The material contained in this publication is considered true and correct as at the date of this publication. The publication is a general guide only prepared solely for the purpose of providing general information in connection with InterGrain, its business and, if applicable the services and products provided by InterGrain. InterGrain does not warrant or guarantee the accuracy, completeness or currency of the publication material and information. InterGrain strongly recommends the publication reader independently research or obtain independent professional advice in connection with the use of the publication material and information for any business decision.

Neither InterGrain, its officers, directors, affiliates or employees, are liable for any cost, expense, damage, liability or loss suffered or incurred by a publication reader or any other party related in any way to the publication reader as a result of the use of publication material and information. Publication Date: October 2026 © Intergrain Pty Ltd 2026. All rights reserved.

Sources
Disease source: 2026 NVT Disease Consensus Ratings. Agronomic characteristics source: InterGrain Breeding.
Yield sourcing: InterGrain long-term MET analysis.